



# **Alaska's Forests** Forested Areas of Alaska 2001 Forests cover 1/3 Of Alaska

## Alaska's Climate...



 Mean annual temperature (MAT) in forested Alaska ranges from ~4°C to -4.9°C

## ...Alaska's Climate



Considerable variation in the length of the growing season throughout Alaska

# **Climate Change**

- Climate related changes have occurred
- During the 20<sup>th</sup> century, boreal Alaska has warmed twice as fast as the global average
- In Interior Alaska:
  - The MAT has increased by 1.3°C over the last 50 years
  - > Precipitation has increased by only 1.4 mm/decade
- Warming since the 1950s appears to be unprecedented in the last 400 years
  - > e.g., Decreased spruce growth in boreal region

# Changes in Disturbance Regimes are Linked to Climate

- i. Wildfire
  - In the last decade, the annual area burned doubled compared to any decade in the last 40 years
- ii. Recent outbreaks of damaging insects (e.g., Dendroctonus rufipennis in the Southcentral and Kenai Peninsula of Alaska)
- iii. The rate of new introductions of exotic flora taxa has increased from 1 to 3 species/year (1941-1968 and 1968-2006, respectively)

# Collaborative Assessment of Climate Change Effects in Alaska's Forests

- In February 2010, university, state and federal scientists met to develop a <u>strategy to assess the</u> <u>impact of climate change</u> on Alaska's forests
- The goals were to:
  - 1) Develop a conceptual framework;
  - 2) Summarize the projected changes in key climate variables;
  - 3) Evaluate the global implications and feedbacks that may alter the rate of changes;
  - 4) Summarize the regional societal consequences;

### Peer-reviewed paper resulting from this collaborative effort



#### **ECOSPHERE**

### Evidence and implications of recent and projected climate change in Alaska's forest ecosystems

Jane M. Wolken, <sup>1,†</sup> Teresa N. Hollingsworth, <sup>2</sup> T. Scott Rupp, <sup>1</sup> F. Stuart Chapin, III, <sup>3</sup> Sarah F. Trainor, <sup>4</sup> Tara M. Barrett, <sup>5</sup> Patrick F. Sullivan, <sup>6</sup> A. David McGuire, <sup>7</sup> Eugenie S. Euskirchen, <sup>3</sup> Paul E. Hennon, <sup>8</sup> Erik A. Beever, <sup>9</sup> Jeff S. Conn, <sup>10</sup> Lisa K. Crone, <sup>11</sup> David V. D'Amore, <sup>8</sup> Nancy Fresco, <sup>1</sup> Thomas A. Hanley, <sup>8</sup> Knut Kielland, <sup>3</sup> James J. Kruse, <sup>12</sup> Trista Patterson, <sup>11</sup> Edward A. G. Schuur, <sup>13</sup> David L. Verbyla, <sup>14</sup> and John Yarie <sup>14</sup>

Scenarios Network for Alaska and Arctic Planning, University of Alaska, 3352 College Road, Fairbanks, Alaska 99709 USA
 <sup>2</sup>United States Department of Agriculture Forest Service, Pacific Northwest Research Station,
 Boreal Ecology Cooperative Research Unit, P.O. Box 756780, University of Alaska, Fairbanks, Alaska 99775 USA
 <sup>3</sup>Institute of Arctic Biology, University of Alaska, Fairbanks, Alaska 99775 USA
 <sup>4</sup>Alaska Center for Climate Assessment and Policy, University of Alaska, 3352 College Road, Fairbanks, Alaska 99709 USA
 <sup>5</sup>United States Department of Agriculture Forest Service, Pacific Northwest Research Station,
 Anchorage Forestry Sciences Laboratory, 3301 C Street, Suite 200, Anchorage, Alaska 99503 USA
 <sup>6</sup>Environment and Natural Resources Institute, Department of Biological Sciences, University of Alaska,
 Anchorage, Alaska 99508 USA

 <sup>7</sup>United States Geological Survey, Alaska Cooperative Fish and Wildlife Research Unit, University of Alaska,

Fairbanks, Alaska 99775 USA

<sup>8</sup>United States Department of Agriculture Forest Service, Pacific Northwest Research Station,
Juneau Forestry Sciences Laboratory, 11305 Glacier Highway, Juneau, Alaska 99801 USA


<sup>9</sup>Northern Rocky Mountain Science Center, United States Geological Survey, Bozeman, Montana 59715 USA

<sup>10</sup>Agricultural Research Service, United States Department of Agriculture, University of Alaska,
Fairbanks, Alaska 99775 USA

<sup>11</sup>United States Department of Agriculture Forest Service, Pacific Northwest Research Station,
 Alaska Wood Utilization Research and Development Center, 204 Siginaka Way, Sitka, Alaska 99835 USA
 <sup>12</sup>Forest Health Protection, State and Private Forestry, United States Department of Agriculture Forest Service,
 Fairbanks Unit, 3700 Airport Way, Fairbanks, Alaska 99709 USA
 <sup>13</sup>Department of Biology, University of Florida, 220 Bartram Hall, P.O. Box 118526, Gainesville, Florida 32611 USA
 <sup>14</sup>Department of Forest Sciences, School of Natural Resources and Agricultural Sciences,
 University of Alaska, Fairbanks, Alaska 99775 USA

Citation: Wolken, J. M., et al. 2011. Evidence and implications of recent and projected climate change in Alaska's forest ecosystems. Ecosphere 2(11):124. doi:10.1890/ES11-00288.1

# The Conceptual Framework



## The Process...

We identified the primary climate drivers, biophysical factors and types of change

#### **Climate Drivers**

= the climate variables (e.g., wind, surface air temperature, precipitation) that directly/ indirectly affect the biophysical subsystem


### **Biophysical Factors**

= the primary categories of disturbances and surface characteristics (e.g., insects, disease, invasive species, permafrost, wildfire) that change in response to the climate drivers

### Types of Change


= the categories of change that occur in all forests (e.g., changes in environment, succession, and biota)

## The Process...






## **Interior-Boreal**



# Interior-boreal: Annual Area Burned



## Interior-Boreal: Wildfire

#### **Changes in Succession**

- Historically, black spruce forests burned during stand replacing fires every 70-130 yrs
  - Low severity wildfires in combination with black spruce traits and understory species led to a resilient forest type
- Post-fire succession has shifted towards deciduous dominated forests
  - Fires burn late into the summer
  - ➤ The deeper depth of burn creates a radically different environment for seedling establishment


#### **Changes in Biota**

- Changes in forest composition will increase moose habitat in the short term
- Wildlife species which prefer distinct forest types are predicted to decrease (e.g., red squirrels and caribou)

# Interior-Boreal: Permafrost-Wildfire Interactions


- Following fire, permafrost degradation/aggradation is determined by the thickness of the SOL, and moss species
- Projected warming and an increase in wildfire will increase permafrost thaw post-fire







## Southcentral- and Kenai-Boreal




# Southcentral/Kenai-Boreal: Insects

#### **Changes in Succession**


- Increasing temperatures decreased the life cycle of spruce bark beetles from 2 years to 1 year
- High spruce mortality
  - Led to very high fuel loads
  - Increased the potential for larger, high severity wildfires


#### **Changes in Biota**


 Invasive plant species may increase with the confluence of increasing human population, wildfire potential and the increased likelihood of invasive plants establishing in recently burned areas



### **Comparison of Alaska's Boreal Forest Regions**








## **SUMMARY**

- The conceptual framework provides a visual tool for resource managers and policy makers
  - > Increase understanding of complex interactions
  - > Identify gaps in our knowledge
- Climate changes have important regional consequences for Alaska residents
- Climate changes also impact the global climate system via effects on carbon and radiation budgets



# **Summary: Southcentral-Boreal**

- We can expect an increase in the frequency and severity of insect outbreaks and associated wildfires
- An increase in invasive plant species establishment is also expected, especially in burned areas

# **Summary: Kenai-Boreal**

- Kenai-boreal = CANARY IN THE COAL MINE
- All of the biophysical factors described in our regional conceptual frameworks are altering the structure and function of forests
- Future climate changes will likely be amplified by the confluence of population growth in this region
- We hypothesize that this is the region where the greatest ecological changes will occur in a relatively short period of time

### **CASE STUDY: Alaskan Boreal Forest**

- Boreal forests in Alaska play an important role in global carbon budgets
- Commercial forestry in Alaska's forests is small-scale relative to other areas
- Alaska may serve as a BASELINE from which to measure our ability to practice natural disturbance-based forest management in other areas

## Acknowledgements

- •Coauthors: T.N. Hollingsworth, T.S. Rupp, S.F. Trainor, T.M. Barrett, E.A. Beever, F.S. Chapin, III., J.S. Conn, L.K. Crone, D.V. D'Amore, E.S. Euskirchen, N. Fresco, T.A. Hanley, P.E. Hennon, K. Kielland, J.J. Kruse, A.D. McGuire, T. Patterson, E.A.G. Schuur, P.F. Sullivan, D.L. Verbyla, and J. Yarie
- •Assistance: Daniel Mann, John Laurence, Tom Kurkowski, Brooke Gamble, and Kimberley Maher
- •Funding: Pacific Northwest Research Station, USDA Forest Service (JVA# 09-JV-11261952-015) and the Bonanza Creek LTER program funded jointly by NSF (grant DEB-0423442) and the USDA Forest Service, Pacific Northwest Research Station (grant PNW01-JV11261952-231)

